Foreword

Teofilo Lee-Chiong, Jr., MD
Consulting Editor

Teofilo Lee-Chiong, Jr., MD
Section of Sleep Medicine
National Jewish Medical and Research Center
University of Colorado Health Sciences Center
1400 Jackson Street
Room J232
Denver, CO 80206, USA

E-mail address:
lee-chiongt@njc.org

A bidirectional relationship exists between the sciences of psychiatry and sleep medicine. Indeed, symptoms of psychiatric disorders are modified by and, more importantly, can lead to sleep disruption. The association of insomnia and the risk of a new psychiatric disorder, specifically major depression, developing is well described. Furthermore, psychiatric disorders can give rise to complaints of insomnia (eg, bipolar disorder, depression, generalized anxiety disorder, obsessive-compulsive disorder, panic disorder, personality disorders, posttraumatic stress disorder, and schizophrenia) or excessive daytime sleepiness (eg, atypical depression and seasonal affective disorder). Certain parasomnias, such as nightmares and sleep terrors, may be more prevalent in patients with psychiatric illnesses. Finally, the medications used to manage psychiatric disorders, including many antidepressant and antipsychotic agents, can affect sleep quality, duration, and architecture.

The clinical course of schizophrenia may be complicated by sleep disturbance, sleep-initiation and sleep-maintenance, insomnia, reversal of day-night sleep patterns, or alternating phases of sleeplessness and sleepiness. Since some antipsychotic

agents can cause sedation, insomnia may also develop following discontinuation of these medications. During exacerbations of psychotic symptoms, prolonged periods of waking may be maintained and terminated only by exhaustion. Conversely, rebound sleepiness can occur during the waning phase of schizophrenia or during residual schizophrenia.

Insomnia is common among persons suffering from mood disorders, and there is generally a correlation between the severity of both conditions. Sleep disturbances and changes in sleep architecture (ie, increase in sleep onset latency or reductions in sleep efficiency, N3 sleep, and REM sleep latency) may both precede or persist after remission of major depressive episodes. Insomnia can be especially severe during a manic episode. Excessive daytime sleepiness, along with an increase in the requirement for sleep, may be seen during the depressive phase of a bipolar disorder, seasonal affective disorder, or atypical depression. In seasonal affective disorder, major depressive episodes occur during the fall and winter, when patients may complain of daytime sleepiness, fatigue, and decreased energy levels; during spring and summer,

some patients' moods improve, but they may experience hypomanic symptoms. Patients with atypical depression may present with lethargy, increase in appetite, weight gain, sensation of heaviness in the extremities, sensitivity to rejection, and excessive sleepiness.

Anxiety disorders, including acute stress disorder, generalized anxiety disorder, and posttraumatic stress disorder, commonly manifest with insomnia, frequent nighttime awakenings, recurring anxiety dreams, or excessive daytime sleepiness. It is important to distinguish generalized anxiety disorder from psychophysiologic insomnia in which anxiety is chiefly restricted to the issues related to sleep disturbance and insomnia. Patients with posttraumatic stress disorder may describe

re-experiences of the original event in frequent distressing dreams, nightmares, and sleep terrors.

In panic disorder, panic attacks can occur during sleep, typically in the transition between light NREM sleep and N3 sleep, but occasionally from REM sleep; awakenings can be accompanied by sympathetic activation and delayed return to sleep. Sleep panic attacks can be triggered by sleep deprivation.

Finally, other psychiatric disorders can also profoundly affect sleep, including insomnia in somatization disorders, obsessive-compulsive personality disorder, and anorexia nervosa; excessive sleepiness in bulimia; and greater rates of obstructive sleep apnea and periodic limb movement disorder in attention deficit hyperactivity disorder.